

# SUMMER ANNUAL GRASS SELECTION

## AS 9301 OR AS 9302 BMR SUDANGRASS

Dry Hay (reccommended to use a mower conditioner)

Haylage Grazing

Seeding Rate: 15-25 lbs/A Seeding Depth: 0.5-0.75"

**Risks:** Prussic acid poisoning; sugarcane aphid pressure;

nitrate poisoning Yield: High

Estimated days to harvest: 40-45

Harvest Height: 24-36" **Residual Height:** 8-10"

Fertilizer Requirement: Moderate

#### PRIME 180 OR PRIME 360 BMR MILLET

Dry Hay (reccommended to use a mower conditioner)

Haylage Grazing

Seeding Rate: 15-20 lbs/A Seeding Depth: 0.25-0.5"

**Risks:** Small seed, tougher to get the correct seed depth;

nitrate poisoning Yield: Moderate

Estimated days to harvest: 40-45

## **KINGFISHER FIBER PRO 70 FORAGE SORGHUM**

Haylage Silage

Seeding Rate: Specific to each variety

Seeding Depth: 1-2"

Risks: Prussic acid poisoning; sugarcane aphid pressure; ni-

trate poisoning **Yield:** High

**Estimated days to harvest:** Specific to each variety

Harvest State: Soft dough or boot stage Residual Height: If regrowth is desired, 8-10"

Fertilizer Requirement: Moderate

### **KINGFISHER SUGAR PRO 55 SORGHUM SUDAN**

Haylage Grazing Silage

Seeding Rate: 40-50 lbs/A Seeding Depth: 1-1.5"

**Risks:** Prussic acid poisoning; sugarcane aphid pressure;

nitrate poisoning Yield: High

Estimated days to harvest: 40-45

Harvest Height: 24-48" Residual Height: 8-10"

## **SOUTHEAST COATED RED RIVER CRABGRASS**

Dry Hay Grazing

Seeding Rate: 5-8 lbs/A

Seeding Depth: Broadcast and Cultipack/Drag or Drill

**Risks:** If allowed to go to seed, will return the following year;

seed to soil contact is critical due to small seed size

**Yield:** Moderate

Estimated days to harvest: 35-40

Harvest Height: 8-12"

#### **MOXIE TEFF**

Dry Hay Grazing

Seeding Rate: 5-8 lbs/A

**Seeding Depth:** Broadcast/Cultipack or Drill 0.25" Risks: Very small seed, establishment is tricky; must be

taken for dry hay in the first cutting

**Yield:** Moderate

Estimated days to harvest: 35-40

Harvest Height: 8-12"



